

Crops(s) **Grape**

Disease(s) **Powdery mildew**

Resistance management strategy for:

Group 3	Demethylation inhibitors (DMI);
Group 5	Amines (morpholines);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7+3	SDHI + DMI
Group 7+12	SDHI + phenylpyrroles (PP);
Group 11	Quinone outside inhibitors (QoI);
Group 11+3	QoI + DMI;
Group 13	Aza-naphthalenes;
Group 13+3	Aza-naphthalenes + DMI;
Group 19	Chitin synthase inhibitor
Group U6	Phenyl-acetamide; and
Group 50	Actin disruptors (aryl-phenyl-ketones).

Guidelines:

1. Apply all these fungicides preventatively.
2. Consecutive applications include from the end of one season to the start of the next. Medium to high risk fungicides (**Group 7** and **11**) if used consecutively should be applied in a mixture or co-formulation with a registered, alternative mode of action for which resistance is not known – where these fungicides have been routinely used for many seasons, field research indicates there is an increased risk of powdery mildew resistance. To ensure effective powdery mildew control in these circumstances, either use alternative modes of action or apply in mixtures.
3. **Do not** apply more than two consecutive sprays of **Group 3, 5, 13, 19, 50** and **U6**.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.

		Group							
		3	5	7 (Incl. 7+3, 7+12)	11 (Incl. 11+3)	13 (Incl. 13+3)	19	50	U6
Total number of powdery mildew targeting sprays	1	1	1	1	1	1	1	1	1
	2	2	1	1	1	2	2	1	1
	3	2	2	1	2	2	2	1	1
	4	2	2	1	2	2	2	2	2
	5	2	2	1	2	2	2	2	2
	6	3	3	2	2	3	3	3	2
	7	3	3	2	2	3	3	3	2
	8	3	3	2	2	3	3	3	2
	9+	3	3	3	2	3	3	3	2

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.