

List of herbicide resistant weeds in Australia

Grass Weeds	Groups	Sites
African lovegrass (<i>Eragrostis curvula</i>)	0 (was J)	8
Annual ryegrass (<i>Lolium rigidum</i>)	1 (was A) 2 (was B) 3 (was D) 5 (was C) 9 (was M) 13 (was Q) 15 (was J, K) 22 (was L) 34 (was Q)	>30,000 >30,000 >5000 >50 >2000 3 >100 >50 3
Annual veldtgrass (<i>Ehrharta longiflora</i>)	1 (was A)	6
Awnless barnyard grass (<i>Echinochloa colona</i>)	9 (was M)	>500
Barnyard grass (<i>Echinochloa crus-galli</i>)	1 (was A) 5 (was C)	1 1
Barley grass (<i>Hordeum</i> spp.)	1 (was A) 2 (was B) 9 (was M) 22 (was L)	>200 >200 8 >100
Brome grass (<i>Bromus</i> spp.)	1 (was A) 2 (was B) 5 (was C) 9 (was M)	>200 >1000 1 5
Large crab grass (<i>Digitaria sanguinalis</i>)	1 (was A) 2 (was B)	2 1
Crowsfoot grass (<i>Eluesine indica</i>)	1 (was A) 22 (was L)	1 5
Feathertop Rhodes grass (<i>Chloris virgata</i>)	9 (was M)	>2000
Giant Parramatta grass (<i>Sporobolus fertilis</i>)	0 (was J)	6
Johnson grass (<i>Sorghum halepense</i>)	9 (was M)	1
Lesser canary grass (<i>Phalaris minor</i>)	1 (was A) 2 (was B)	20 10
Liverseed grass (<i>Urochloa panicoides</i>)	5 (was C) 9 (was M)	7 4
Paradoxa grass (<i>Phalaris paradoxa</i>)	1 (was A) 2 (was B)	7 4
Serrated tussock (<i>Nassella trichotoma</i>)	0 (was J)	32
Silver grass (<i>Vulpia bromoides</i>)	5 (was C)	3

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.

	22 (was L)	1
Sweet summer grass (<i>Brachiaria eruciformis</i>)	9 (was M)	15
Wild oat (<i>Avena</i> spp.)	1 (was A) 2 (was B) 9 (was M) 0 (was Z)	>5000 >200 2 >200
Windmill grass (<i>Chloris truncata</i>)	9 (was M)	13
Winter grass (<i>Poa annua</i>)	1 (was A) 2 (was B) 3 (was D) 5 (was C) 9 (was M) 15 (was J) 31 (was Z)	3 20 >100 10 10 10 3
Broadleaf weeds	Groups	Sites
African turnip weed (<i>Sisymbrium thellungii</i>)	2 (was B)	2
Arrowhead (<i>Sagittaria montevidensis</i>)	2 (was B)	20
Black bindweed (<i>Fallopia convolvulus</i>)	2 (was B)	2
Black nightshade (<i>Solanum nigrum</i>)	22 (was L)	2
Bedstraw (<i>Galium tricornutum</i>)	2 (was B)	3
Calomba daisy (<i>Pentzia suffruticosa</i>)	2 (was B)	2
Capeweed (<i>Arctotheca calendula</i>)	4 (was I) 9 (was M) 12 (was F) 22 (was L)	3 1 1 1
Charlock (<i>Sinapis arvensis</i>)	2 (was B)	2
Common sowthistle (<i>Sonchus oleraceus</i>)	2 (was B) 4 (was I) 9 (was M)	>10,000 >100 >100
Dense-flowered fumitory (<i>Fumaria densiflora</i>)	3 (was D)	2
Dirty dora (<i>Cyperus difformis</i>)	2 (was B)	>50
Iceplant (<i>Mesembryanthemum crystallinum</i>)	2 (was B)	2
Fleabane (<i>Conyza bonariensis</i>)	2 (was B) 22 (was L) 9 (was M)	>100 1 >2000
Indian hedge mustard (<i>Sisymbrium orientale</i>)	2 (was B) 4 (was I) 5 (was C) 12 (was F)	>1000 >50 16 >50

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.

Lincoln weed (<i>Diploaxis tenuifolia</i>)	2 (was B)	20
Paterson's curse (<i>Echium plantagineum</i>)	2 (was B)	2
	9 (was M)	1
Pennsylvania cudweed (<i>Gamochaeta pensylvanica</i>)	22 (was L)	2
Prickly lettuce (<i>Lactuca serriola</i>)	2 (was B)	>2000
	9 (was M)	1
Small square weed (<i>Mitracarpus hirtus</i>)	22 (was L)	1
Starfruit (<i>Damasonium minus</i>)	2 (was B)	5
Stinging nettle (<i>Urtica urens</i>)	5 (was C)	1
Tall fleabane (<i>Conyza sumatrensis</i>)	9 (was M)	10
Tridax (<i>Tridax procumbens</i>)	9 (was M)	1
Turnip weed (<i>Rapistrum rugosum</i>)	2 (was B)	5
Wild radish (<i>Raphanus raphanistrum</i>)	2 (was B)	>5000
	4 (was I)	>1000
	5 (was C)	>20
	9 (was M)	3
	12 (was F)	>1000
	27 (was H)	1
	29 (was O)	2
Wild turnip (<i>Brassica tournefortii</i>)	2 (was B)	>100
Willow-leaved lettuce (<i>Lactuca saligna</i>)	9 (was M)	2
Winged Slender Thistle (<i>Carduus tenuiflorus</i>)	4 (was I)	2

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.