

Crops(s) **Wheat**

Disease(s) **Yellow spot**

Resistance management strategy for:

Group 3	Demethylation inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7+3	SDHI + DMI;
Group 11+3	Quinone outside inhibitors (QoI) + DMI;
Group 11+4	QoI + phenylamides (PA);
Group 11+7+3	QoI + SDHI + DMI; and
Group 50*	Actin disruption aryl-phenyl-ketones.

Guidelines:

1. Fungicides should be used as protectant treatments prior to symptoms of disease when conditions are conducive to disease development.
2. In high-risk disease environments, integrated management approaches should be used to reduce fungicide resistance risk, which may include:
 - a. removal or burning of stubble,
 - b. crop rotation (**avoid** wheat on wheat),
 - c. control of green bridge volunteers; and
 - d. use of tolerant and resistant varieties.
3. **Do not** apply more than two applications per growing season of **Group 11**, or **7** containing products. This includes in-furrow or seed treatments that have activity on foliar diseases. Combinations of in-furrow and seed treatment are counted as one application.
4. **Do not** apply consecutive applications of **Group 11** containing products. This includes in-furrow i.e. If a **Group 11+4** fungicide has been used in-furrow at planting, the first foliar fungicide spray must not contain a **Group 11** fungicide.
5. If a **Group 7** seed treatment has been used with foliar activity (as determined by label claims), the first foliar fungicide applied should not contain a **Group 7** fungicide.
6. **Do not** apply more than one application of a **Group 7** seed treatment with foliar activity in any two consecutive growing seasons.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.

7. If a **Group 7** or **11** fungicide is being applied as a foliar spray, it must be in a co-formulation or in mixture with a registered mixing partner with a different mode of action, with no known resistance. The mixture partner should also provide effective disease control when used alone on the target disease.
8. **Do not** apply more than three applications containing **Group 3** fungicides per growing season. This total of three applications includes DMIs applied as **Group 11+3** or **Group 11+7+3** co-formulations and in-furrow or seed treatments that have activity on foliar diseases. Combinations of in furrow and seed treatment are counted as one application.
9. Minimise use of **Group 3** fungicides which are known to have compromised resistance status.
10. **Do not** apply **Group 13** fungicides onto existing infections of powdery mildew. Always use preventatively. If applying consecutive applications, use an effective mixing partner.
11. Reduced rates, including multiple applications must not be used. Always use label rates.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.