

**Crops(s)**            **Almond**

**Disease(s)**        **Rust**

Resistance management strategy for:

<b>Group 3</b>	Demethylation inhibitors (DMI);
<b>Group 7</b>	Succinate dehydrogenase inhibitors (SDHI);
<b>Group 7 + 3</b>	SDHI + DMI;
<b>Group 11</b>	Quinone outside inhibitors (QoI);
<b>Group 11+3</b>	QoI + DMI; and
<b>Group 11+7</b>	QoI + SDHI.

		Maximum recommended number of sprays which can contain		
		Group:		
		<b>3</b>	<b>7</b> (incl. 7+11 and 7+3)	<b>11</b> (incl. 11+3 and 7+11)
Total number of rust targeting sprays	<b>1</b>	1	1	1
	<b>2</b>	2	1	1
	<b>3</b>	2	1	2
	<b>4</b>	2	1	2
	<b>5</b>	2	1	2
	<b>6</b>	3	2	2
	<b>7</b>	3	2	2
	<b>8</b>	3	2	2
	<b>9+</b>	3	3	3

**Guidelines:**

1. **Do not** apply consecutive sprays of products containing **Group 7** or **11** if applied solo. Consecutive sprays include mixtures containing **Group 7** or **11**.
2. If applying **Group 7** or **Group 11** fungicides in mixtures e.g. **Group 11+7**, consecutive application is allowed. **Do not** apply more than two consecutive sprays before changing to another group.

**Please note:**

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.

- 
3. If two consecutive sprays of **Group 3** or **Group 11+3** fungicides are used, then use the same number of sprays of an alternative group(s) before using another **Group 3**, including sprays in the following seasons.
  4. Rotate with products from **Group M**.
  5. No specific resistance management strategy has been developed for low-risk fungicides, including those in **Group M** and **BM**. These products should be included in a management strategy as per label recommendations.
  6. The spray program should be considered and the strategy applied on a whole-orchard basis.

---

**Please note:**

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.

