

Crops(s) **Various**

Insect(s) **Mites**

Guidelines:

1. For [Pome fruits](#), [Strawberries/ornamentals](#) and [Redlegged earth mite](#) see specific CropLife strategies.
2. Monitor mite activity and treat infestations before thresholds are reached, i.e. spray earlier rather than later. Seek advice on local threshold levels.
3. **Do not** apply sequential applications of products from any one chemical group.
4. Preferably products with the same Mode of Action should not be used more than twice in a growing season.
5. Incorporate the use of predatory mites for the control of this pest wherever possible.

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a. Product labels should at all times be carefully read and adhered to;
- b. Full recommended rates of registered insecticides should always be used; and
- c. Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.